

| Public Presentation at the Faculty of Physics Date: 25 February 2020                  |                                   |  |                                    |
|---|-----------------------------------|--|------------------------------------|
| Place: Josef-Stefan lecture hall, 3 <sup>rd</sup> floor, Boltzmannngasse 5, 1090 Wien |                                   |  |                                    |
| Time Slot   | Name                              | Topic  | Supervisor                         |
| 9:00-9:10   | --                                | Introduction by Bogdan Sepiol  |                                    |
| 9:10-9:25   | Rieser Philipp                    | Time domain interference of biomolecules and quantum assisted spectroscopy   | Markus Arndt                       |
| 9:25-9:40   | Troyer Stephan                    | Levitated optomechanics with aspheric nanoparticles and microcavities  | Markus Arndt                       |
| 9:40-9:55   | Lanz Andreas                      | Physics of a non-neutral plasma within a cusped magnetic field   | Eberhard Widmann                   |
| 9:55-10:10  | Nowak Lilian                      | Deexcitation and cooling techniques for precision measurements with antihydrogen   | Eberhard Widmann                   |
| <b>10:10-10:40</b>  | <b>Bernd Aichner</b>              | <b>Tuning the electronic properties of superconductors by Helium-ion-irradiation</b>                                       |                                    |
| 10:40-11:10   | COFFEE BREAK                      |  |                                    |
| 11:10-11:25   | Tüchler Marlene                   | Probing the strong interaction with kaonic atom x-ray measurements at low energies   | Eberhard Widmann<br>Johann Zmeskal |
| 11:25-11:40   | Postl Andreas                     | Automated manipulation of single atoms using focused electron irradiation  | Toma Susi                          |
| 11:40-11:55   | Barzegar Hamed                    | Cosmological non-vacuum spacetimes   | Fajman David                       |
| 11:55-12:10   | Fiore Mosca Dario                 | Quantum Magnetism in Relativistic Oxides   | Cesare Franchini                   |
| <b>12:10-12:40</b>  | <b>Uros Delic</b>                 |  |                                    |
| 12:40-13:40   | LUNCH BREAK                       |  |                                    |
| 13:40-13:55   | Kysela Jaroslav                   | Entanglement, interference and manipulation of systems in high dimensions  | Anton Zeilinger                    |
| 13:55-14:10   | Ecker Sebastian                   | Entanglement distribution over noisy quantum channels  | Anton Zeilinger                    |
| 14:10-14:25   | Rosenberg Margaret                | Phase behavior of colloidal magnetic paterlets   | Sofia Kantorovich                  |
| 14:25-14:40   | Mostarac Deniz                    | Structural and magnetic properties of supracolloidal, nanoscopic magnetic filaments and filament based soft matter systems | Sofia Kantorovich                  |
| 14:40-15:10   | <b>Iurii Chubak</b>               | <b>Active topological glass</b>  |                                    |
| 15:10-15:30   | COFFEE BREAK                      |  |                                    |
| 15:30-15:45   | Sukurma Zoran                     | Auxiliary field quantum Monte Carlo method in projector augmented-wave method  | Georg Kresse                       |
| 15:45-16:00   | Maestre Vazquez Dante             | Enhancing Phase Imaging with Local Wavefront Shaping Techniques  | Thomas Juffmann                    |
| 16:00-16:15   | Wirtitsch Daniel                  | Quantum Sensing using the Nitrogen Vacancy Center in Diamond   | Philip Walther                     |
| 16:15-16:30   | Silvestri Raffaele                | Gravitational induced phase shift on single photons  | Philip Walther                     |
| 16:30-16:45   | Trillo Fernandez David            | Exploring time in non-relativistic quantum mechanics   | Brukner Caslav                     |
| 16:45-17:00   | Chirita Mihaila Marius Constantin | Wavefront shaping of free electrons using light  | Thomas Juffmann                    |

# Tuning the electronic properties of superconductors by Helium-ion-irradiation

Bernd Aichner<sup>1</sup>, Benedikt Müller<sup>2</sup>, Kristijan Luka Mletschnig<sup>1</sup>, Max Karrer<sup>2</sup>, Johannes David Pedarnig<sup>3</sup>, Vyacheslav R. Misko<sup>4</sup>, Dieter Kölle<sup>2</sup>, and Wolfgang Lang<sup>1</sup>

<sup>1</sup>University of Vienna

<sup>2</sup>University of Tübingen


<sup>3</sup>Johannes Kepler University Linz

<sup>4</sup>University of Antwerpen

Superconductors exhibit intriguing electronic properties when they are cooled below their critical temperature  $T_c$ , but their application is by now limited because of the high demand on cooling power. One way out could be the use of high- $T_c$ -superconductors as for example  $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$  (YBCO) which has a critical temperature above the boiling point of liquid nitrogen and allows thus for relatively cheap operation. One drawback in the fabrication of electronic devices from this material is its sensitivity to surface damage, which limits the use of conventional etching techniques. A promising method to realize nano-structures in such materials is the local irradiation with light ions, which makes the material non-superconducting instead of removing it.

This can be done, for example, by exposing the superconducting film to the collinear beam of an ion implanter and shading the parts which are intended to remain superconducting by a mask. Indeed, many experiments have been successfully conducted with this method and it was shown that it is suitable to produce structures with a size of a few hundreds of nanometers [1]. However, we needed to find another way to produce smaller structures and found it in the use of a helium-ion microscope. These machines have been developed only recently (For further information, see for example ref. [2]) and allow for surface-sensitive imaging with a resolution of 0.5 nm. By intentionally defocusing the beam, they are also suitable to produce nano-structures in thin films of high-temperature superconductors. We use this technique to create artificial defect columns in YBCO films, and investigate the resulting electronic properties by electronic transport measurements at temperatures between 4.2 K and 300 K and in magnetic fields up to 1 T. Since magnetic fields penetrate into a type-II superconductor like YBCO in the form of single flux quanta, the motion of these flux quanta can be used as a method to investigate the influence of the artificially created defect columns. The flux quanta are preferentially anchored in defects and since their motion causes dissipation, stable arrangements that impede motion lead to high critical currents and low resistance.

The aim of this talk is to introduce the concept of manipulating flux quanta and tuning thus the electronic transport properties of YBCO films by the introduction of artificial defect columns. The design of a helium-ion microscope is briefly described and its applicability to produce defects in superconductors is discussed. The results of electronic transport measurements are presented, which show the practicability of this method [3] and the possibility to switch between two stable states of flux quanta arrangements in a superconducting structure realized by this technique [4].



## References

- [1] V. Moshchalkov, R. Wördenweber, and W. Lang, *Nanoscience and Engineering in Superconductivity*. Springer Berlin Heidelberg, 2010.
- [2] G. Hlawacek and A. Götzhäuser, *Helium Ion Microscopy*. Springer International Publishing: Switzerland, 2016.
- [3] B. Aichner, B. Müller, M. Karrer, V. R. Misko, F. Limberger, K. L. Mletschnig, M. Dosmailov, J. D. Pedarnig, F. Nori, R. Kleiner, D. Koelle, and W. Lang, “Ultradense Tailored Vortex Pinning Arrays in Superconducting  $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$  Thin Films Created by Focused He Ion Beam Irradiation for Fluxonics Applications,” *ACS Applied Nano Materials*, vol. 2, pp. 5108–5115, Jul 2019.
- [4] B. Aichner, K. L. Mletschnig, B. Müller, M. Karrer, M. Dosmailov, J. D. Pedarnig, R. Kleiner, D. Koelle, and W. Lang, “Angular magnetic-field dependence of vortex matching in pinning lattices fabricated by focused or masked helium ion beam irradiation of superconducting  $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$  thin films,” *Low Temperature Physics/Fizika Nizkikh Temperatur*, vol. 46, no. 4, 2020. in press (to appear on February 28, 2020).

# Active topological glass

Iurii Chubak

The glass transition in soft matter systems is generally triggered by an increase in packing fraction or a decrease in temperature. It has been conjectured that the internal topology of the constituent particles, such as polymers, can cause glassiness too.

However, the conjecture relies on immobilizing a fraction of the particles and is therefore difficult to fulfill experimentally. Here we show that in dense solutions of circular polymers containing (active) segments of increased mobility, the interplay of the activity and the topology of the polymers generates an unprecedented glassy state of matter. The active isotropic driving enhances mutual ring threading to the extent that the rings can relax only in a cooperative way, which dramatically increases relaxation times.

